



NAPAN© Standards for Practice, 3rd Edition, 2014

EDITOR'S ERRATA

Please note the following typographical errors in this document:

1. Resource 11: Management of Postoperative Nausea and Vomiting in all PeriAnesthesia Phases

The following statement, 2.5.6.1iii. on P. 198 (English):

“**2.5.6.1iii.** Health Canada has since allowed limited use of the butyrophenones to those who are not predisposed to prolonged QT intervals (greater than **.044**), electrolyte imbalance.....”

Should read,

“ **2.5.6.1iii.** Health Canada has since allowed limited use of the butyrophenones to those who are not predisposed to prolonged QT intervals (greater than **0.44**), electrolyte imbalance.....”

2. Resource 11: Management of Postoperative Nausea and Vomiting in all PeriAnesthesia Phases

The following statement, 2.5.7 on P. 199 (English):

“**2.5.7** Dimenhydrinate and *dimenhydramine* are effective on the histamine and dopamine receptors and are effective for clients undergoing surgery affecting the vestibular apparatus”

Should read,

“**2.5.7** Dimenhydrinate and *diphenhydramine* are effective on the histamine and dopamine receptors and are effective for clients undergoing surgery affecting the vestibular apparatus”

3. Resource 13: Airway Management in PostAnesthesia Phase I:

The following statement, 5.2.3.6, on P. 303 (English):

“**5.2.3.6** Just prior to withdrawal of the ETT, instruct the client to inhale deeply and then exhale, removing the tube on **end-expiration**”

Should read,

“**5.2.3.6** Just prior to withdrawal of the ETT, instruct the client to inhale deeply and then exhale, removing the tube on **end-inspiration (pre-expiration)**”

We apologize for any inconvenience this may cause the reader.

NAPAN© Board of Directors, 2014